BEETLES (Coleoptera)

Isually hard-bodied insects, with FW thickened and meeting in a traight line down back; antennae nearly always with 10 or more egments; mouth parts chewing.

HECKERED BEETLES, Family Cleridae p. 165 Often marked with red, orange, yellow, or blue; pronotum narrower than FW; body and legs with erect pubescence; tarsi 5-5-5.

Enoclerus ichneumoneus (Fabricius); e. U.S.

ERMESTID BEETLES, Family Dermestidae p. 164 Elongate to broadly oval, often with scales or hair forming patterns; antennae short and clubbed; tarsi 5-5-5.

CARPET BEETLE, Anthrenus scrophulariae (Linn.); N. Amer. OFT-WINGED FLOWER BEETLES, Family Malachiidae p. 164 Body soft and wedge-shaped, FW broad toward rear; antennae serrate; black, blue, or green, often with orange, red, or yellow; tarsi 5-5-5; abdomen with 6 or fewer segments. Collops vittatus (Say); w. U.S.

IGHTNINGBUGS or FIREFLIES, Family Lampyridae p. 162 Elongate, parallel-sided, rounded at ends; soft-bodied; head concealed by pronotum; abdomen often luminescent. Photuris pennsylvanica (De Geer); e. U.S.

Generally black and shiny, sometimes brightly colored; legs long and slender; head and eyes nearly always narrower than pronotum; tarsi 5-5-5; antennae threadlike.

Calosoma scrutator (Fabricius); e. U.S.

OLDIER BEETLES, Family Cantharidae p. 162 Similar to lightningbugs (above), but head not concealed and no abdominal segments luminescent; black or brown, often with red, yellow, or orange, some mostly yellowish. Chauliognathus pennsylvanicus De Geer; e. U.S.

IET-WINGED BEETLES, Family Lycidae p. 163 FW with a network of ridges; body usually broadest toward rear; head concealed; black, some with red or yellow. Calopteron terminale (Say); U.S.

NETALLIC WOOD-BORING BEETLES,

p. 169

amily Buprestidae Elongate, usually parallel-sided; body hard, generally metallic, especially below; antennae short, threadlike or serrate; tarsi 5-5-5.

Acmaeodera pulchella (Herbst); e. U.S.

ARRION BEETLES, Family Silphidae p. 158
Black, often with yellow, orange, or red; antennae clubbed; tarsi
5-5-5; body flattened; FW sometimes short, exposing 1-3 abdominal segments.

Nicrophorus marginatus Fabricius; N. America.

